

## Original article

# Insecticide resistance studies on German cockroach (*Blattella germanica*) strains to malathion, propoxur and lambda-cyhalothrin

Abed Kakeh-Khani<sup>a</sup>, Mansour Nazari<sup>a</sup>, Hassan Nasirian<sup>b,\*</sup>

<sup>a</sup>Department of Medical Entomology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Islamic Republic of Iran

<sup>b</sup>Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran

---

**Background:** The German cockroach has been known as the most invading human dwelling cockroach species. For managing the insecticide resistance, it is periodically a substantial need to evaluate susceptibility level of the German cockroaches to consuming insecticides.

**Objective:** To evaluate resistance of the German cockroaches to malathion, propoxur and lambda-cyhalothrin.

**Methods:** Three cockroach strains including susceptible reference, and two wild strains were provided, maintained and colonized. The wild strains were collected from two hospitals of Sanandaj in Iran. Surface contact method was used for bioassay using standard glass jar procedure. The adult male cockroaches were treated with 5 - 6 insecticide dose exposures at a 30-min period. The dose exposures of malathion were 6.25, 12.5, 25, 50 and 100, and 25, 50, 100, 200 and 400 mg/m<sup>2</sup> for susceptible and wild cockroach strains, respectively. The dose exposures of propoxur were 5, 10, 20, 40 and 80, and 20, 40, 80, 160 and 320 mg/m<sup>2</sup> for susceptible and wild cockroach strains, respectively. The dose exposures of lambda-cyhalothrin were 0.075, 0.15, 0.3, 0.6 and 1.2, and 0.3, 0.6, 1.2, 2.4 and 4.8 mg/m<sup>2</sup> for susceptible and wild cockroach strains, respectively. Each dose exposures were replicated 3 times (10 cockroaches for each replicate). Control groups received acetone alone.

**Results:** The lethal doses<sub>50</sub> (LDs<sub>50</sub>) of the susceptible strain were 30.01, 16.56 and 0.303 for malathion, propoxur and lambda-cyhalothrin respectively. While the LDs<sub>50</sub> of the wild cockroach strains were 157.45, 74.59 and 0.506 for malathion, propoxur and lambda-cyhalothrin, respectively. Compared to the susceptible strain, the wild German cockroach strains were more tolerant to malathion and propoxur. Compared to the susceptible strain the wild German cockroach strains were more sensitive to lambda-cyhalothrin, confirmed also by a significant difference ( $P = 0.033$ ). The results of probit analysis and regression lines of treated insecticides indicate that the German cockroach was resistant to malathion and propoxur while susceptible to lambda-cyhalothrin. The resistance ratio<sub>50</sub> (RR<sub>50</sub>) of the malathion and propoxur insecticides were about 5.0 folds.

**Conclusion:** The study supports that malathion and propoxur should not be used for control of the German cockroaches. An appropriate choice is to use lambda-cyhalothrin due to negative cross-resistance effect which occurs here. The study also provides a baseline of resistance data towards a few insecticides in a few German cockroach populations in Iran.

**Keywords:** *Blattella germanica*, insecticide resistance, lambda-cyhalothrin, malathion, propoxur.

---

Among cockroaches of various species, the German cockroach (*Blattella germanica*: Blattaria,

Blattellidae) has been known as the most invading cockroach species of the human dwelling areas<sup>(1-9)</sup> with a variety of medical importance.<sup>(9-15)</sup> Recently, it has been observed that about 25 fungal and 61 bacterial species were recovered from the German cockroaches associated with high morbidity and mortality in immune-compromised patients in the hospital environments.<sup>(16-17)</sup> Therefore the control of this important human dwelling pest is necessary. The use of insecticides is still the most important tool

---

\*Correspondence to: Hassan Nasirian, Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran.

E-mail: hanasirian@yahoo.com

Received: January 30, 2020

Revised: March 13, 2020

Accepted: April 8, 2020

for control of the German cockroaches.<sup>(18)</sup> In order to predict and manage the insecticides resistance phenomenon, it is periodically a substantial need to evaluate the sensitivity level of the German cockroach to consuming insecticides. Various studies have been done in this area<sup>(19 - 23)</sup> but there is a need to be performed in other region. In this regard, a study is designed to evaluate the sensitivity level of the German cockroaches to malathion, propoxur and lambda-cyhalothrin insecticides. The aim of this study is to determine the susceptibility level of the wild German cockroach strains from Sanandaj of Iran to malathion, propoxur and lambda-cyhalothrin insecticides.

## Materials and methods

### *Sites of cockroach collection*

Cockroach collecting was done in Sanandaj, the center of Kurdistan Province in 2016. Kurdistan Province is home to natural beauty. It has abundant tourist attractions in terms of cultural heritage, history and handicrafts. The area has a semi-arid Mediterranean climate and its coordinates are: 35° 20'N - 46° 50'E. To do the study, we provided 3 cockroach strains including a standard susceptible strain, and two wild German cockroach strains. The standard susceptible (SS) strain was provided by the School of Public Health, Tehran University of Medical Sciences (TUMS) which had been maintained since 1975 in the insectary, without exposure to any insecticide. Two wild German cockroach strains including a strain from Besat Hospital of Kurdistan University of Medical Sciences and a strain from a hospital of Iranian Social Security Organization of Kurdistan (Tamin-e Ejtemaei hospital) were collected.

### *Cockroaches collecting and rearing*

Cockroaches were collected using Nasirian H, *et al.* procedure.<sup>(24)</sup> Each cockroach strain after collecting were delivered into a large glass jar, maintained and colonized at 24 ± 2°C, 40 ± 5% RH, and a photoperiod of 12:12 (L:D) h in the insectary at the Department of Medical Entomology, School of Medicine, Hamadan University of Medical Sciences. The cockroaches were provided with rodent diet, a cotton plugged water vial and a cardboard as a harborage.

### *Chemicals and reagents*

Malathion (technical grade, 98.0%), propoxur

(technical grade, 95.0%) and lambda-cyhalothrin (technical grade, 95.0%) as insecticides, CO<sub>2</sub> as an anesthetic, and acetone as a solvent were used. Malathion was provided by Asia Agro Chemicals & Fertilizers (Khurja, Dist. Bulandshahr, India). Propoxur and lambda-cyhalothrin were provided by Kalyani Industries Pvt. Ltd. (Maharashtra, India).

### *Surface contact method bioassay*

Surface contact method was used for bioassay using a glass jar. The glass jar inner surface areas were 188.4 cm<sup>2</sup>. Technical grade of the insecticides was diluted by acetone solvent. To dilute insecticide concentrations 1 ml volume solvent considered contains insecticide doses. Then 1 ml of diluted insecticide was pipetted into the glass jar. To deposit the insecticide evenly over the inner surface of the glass jars, the glass jars were rolled horizontally over a flat surface until all of the acetone had evaporated.

In a series of contact experiments, a concentration of technical grade at a 30-min exposure time was found to be a sufficient time for exposure of adult males to insecticide treated of the glass jar inner surfaces.<sup>(9, 25 - 28)</sup> Then adult male cockroaches were treated with 5 - 6 insecticide dose exposures at a 30-min time. After preliminary experiments, the dose exposures of insecticides were determined. The determined dose exposures of malathion insecticide were 6.25, 12.5, 25, 50 and 100, and 25, 50, 100, 200 and 400 mg/m<sup>2</sup> for susceptible and wild cockroach strains, respectively. Finally, the determined dose exposures of propoxur insecticide were also 5, 10, 20, 40 and 80, and 20, 40, 80, 160 and 320 mg/m<sup>2</sup> for susceptible and wild cockroach strains, respectively. Finally the determined dose exposures of lambda-cyhalothrin insecticide were 0.075, 0.15, 0.3, 0.6 and 1.2, and 0.3, 0.6, 1.2, 2.4 and 4.8 mg/m<sup>2</sup> for susceptible and wild cockroach strains, respectively. Each dose exposures were replicated 3 times (10 cockroaches for each replicate). Control groups received acetone alone. All insecticide dose exposures were given > 0.0% and < 100.0% mortality at 24 h after insecticide dose exposures. Insecticide-exposed males were placed in plastic dishes, provided with food and water, and monitored for mortality for 24 h under the same temperature and photoperiod as the colony. If insects on their backs were unable to right themselves, they were considered dead. All experiments were replicated 3 times and 10 cockroaches for each replicate.

**Statistical analysis**


IBM SPSS Statistics Data Editor Version 23 was used to analyze statistical analysis of the data on a computer. Mortality data from the replicates were pooled and the dose exposure mortality was assessed by probit regression analysis. One-sample Kolmogorov-Smirnov test was used to check the normality of the data distribution. After checking, to make sure that the distribution of data was normal, ANOVA was used to compare treated male cockroaches to insecticides for any significant difference. Microsoft Office Excel (2010 version) was used to calculate equation and draw the mortality and regression lines of insecticide treated cockroaches.

**Results**

**Cockroach insecticide susceptibility level**

Compared to the susceptible strain the wild German cockroach strains were more tolerant to

malathion and propoxur, while they were susceptible to lambdacyhalothrin. In other words, the wild German cockroach strains were resistant to malathion and propoxur. Compared to the susceptible strain, the malathion mortality lines of the wild German cockroach strains locate the right of the susceptible strain. Unlike malathion, the lambdacyhalothrin mortality line of the susceptible strain locate the right of the wild German cockroach strains. While the propoxur mortality lines of the wild and susceptible cockroach strains overlap (Figure 1). Statistical analysis showed a significant difference ( $P = 0.036$ ) between treated insecticides against the German cockroach strains. There was also a significant differences between treated malathion and lambdacyhalothrin ( $P = 0.033$ ) against the German cockroach strains followed by Post Hoc Tests.



**Figure 1.** Mortality of the German cockroach strains after treated insecticides. There is a significant difference ( $P = 0.036$ ) between treated insecticides against the German cockroach strains. There is also a significant difference between treated malathion and lambdacyhalothrin ( $P = 0.033$ ) against the German cockroach strains followed by Post Hoc Tests.

**Table 1.** Probit regression analysis of insecticides against the German cockroach strains.


Strain	n <sup>a</sup>	Intercept ± SE	Slope ± SE	X <sup>2</sup>	df	P	LD <sub>50</sub> <sup>b</sup> (LB-UB) <sup>c</sup>	RR <sub>S0</sub>	LD <sub>90</sub> <sup>b</sup> (LB-UB) <sup>c</sup>	RR <sub>90</sub>
Malathion										
S	150	0.02 ± 0.004	-0.62 ± 0.17	7.4	3	0.06	30.01 (5.7 - 86.5)	0.0	92.57 (57.3 - 814.0)	0.0
BH	150	0.005 ± 0.001	-0.82 ± 0.17	4.7	3	0.19	150.42 (109.3 - 195.1)	5.0	384.24 (312.6 - 519.6)	4.2
TEH	150	0.006 ± 0.001	-0.95 ± 0.18	4.7	3	0.19	164.47 (125.6 - 209.2)	5.5	387.24 (317.8 - 514.5)	4.2
Propoxur										
S	150	0.03 ± 0.006	-0.48 ± 0.17	5.8	3	0.12	16.56 (3.3 - 38.3)	0.0	60.60 (38.7 - 269.2)	0.0
BH	150	0.007 ± 0.001	-0.45 ± 0.17	1.8	3	0.62	67.20 (24.9 - 100.2)	4.1	258.25 (204.3 - 369.9)	4.3
TEH	150	0.006 ± 0.001	-0.49 ± 0.16	5.1	3	0.16	81.98 (37.7 - 118.8)	5.0	297.15 (234.7 - 427.6)	4.9
Lambda-cyhalothrin										
S	150	1.8 ± 0.33	-0.55 ± 0.17	6.6	3	0.08	0.303 (0.07 - 0.75)	0.0	1.006 (0.63 - 5.7)	0.0
BH	150	0.48 ± 0.105	-0.23 ± 0.17	2.3	3	0.51	0.477 (0.05 - 0.97)	1.6	3.15 (2.4 - 4.7)	3.1
TEH	150	0.42 ± 0.009	-0.22 ± 0.17	3.9	3	0.27	0.534 (0.17 - 1.1)	1.8	3.61 (2.8 - 5.4)	3.6

<sup>a</sup>Numbers of cockroach treated; <sup>b</sup>LD<sub>50</sub>, LD<sub>90</sub> values in mg/m<sup>2</sup> (LB, Lower bound and UB, Upper bound of 95% confidence interval); S, Susceptible; BH, Besat hospital and TEH, Tamin-e Ejtemaei hospital strains.

**Probit regression analysis**

The slope of regression lines shows the heterogeneity and genetic variation of a strain population. The higher slope of a regression line indicates the less uniformity of a strain population.

Further analysis also indicates genetic diversity of a strain population. Compared to the susceptible strain the wild German cockroach strains were heterogenous, based on the slope of the regression lines and  $X^2$  of the cockroach population strains (Table 1).



**Figure 2.** Regression lines of treated insecticides against the German cockroach strains.

The lethal doses<sub>50</sub> (LDs<sub>50</sub>) of the susceptible strain were 30.01, 16.56 and 0.303 for malathion, propoxur and lambda-cyhalothrin, respectively. While the LDs<sub>50</sub> of the wild cockroach strains were 157.45, 74.59 and 0.506 for malathion, propoxur and lambda-cyhalothrin, respectively. The resistance ratio<sub>50</sub> (RR<sub>50</sub>) of the malathion and propoxur were about 5.0 folds. While the RR<sub>50</sub> of the lambda-cyhalothrin insecticides were about 1.7 folds. Compared to the susceptible strain, the wild cockroach strains were resistant to malathion and propoxur. Unlike malathion and propoxur, the wild cockroach strains were susceptible to the lambda-cyhalothrin (Table 1). There was also a significant difference between malathion and lambda-cyhalothrin insecticides ( $P = 0.033$ ) against the German cockroach strains.

Compared to the susceptible strain, the regression lines of the wild cockroach strains treated with malathion locate the right of the susceptible strain. Unlike malathion, the lambda-cyhalothrin regression line of the susceptible strain locate the right of the wild cockroach strains, while the propoxur regression lines of the wild and susceptible cockroach strains overlap (Figure 2). Based on the slope of regression lines of the German cockroach strains which were decreased (Table 1), and mortality of the German cockroach strains (Figure 2) and regression lines of treated insecticides (Figure 2), the population of the wild German cockroach strains were heterogeneous.

## Discussion

The present study was designed to evaluate the susceptibility level of the German cockroach strains to malathion, propoxur and lambda-cyhalothrin insecticides for managing the insecticide resistance.

According to probit analysis and regression lines of the treated insecticides, the results of the study indicate that the German cockroach was resistant to malathion and propoxur while susceptible to lambda-cyhalothrin. The RR<sub>50</sub> of the malathion and propoxur insecticides were about 5.0 folds and the malathion regression lines locate the right of the susceptible strain. Further analysis showed a significant difference between treated insecticides against the German cockroach strains. There was a significant difference between treated malathion and lambda-cyhalothrin insecticides against the German cockroach strains followed by Post Hoc Tests.

In accord with this study, it is well documented that the German cockroach is resistant to

malathion<sup>(21,29-33)</sup> and propoxur.<sup>(29-40)</sup> While unlike the current study, it is also well documented that the German cockroach is resistant to lambda-cyhalothrin.<sup>(32,33,41-43)</sup> Unlike malathion and propoxur insecticides, the wild cockroach strains were susceptible to lambda-cyhalothrin. Compared to the susceptible strain the lambda-cyhalothrin mortality and regression lines of the wild cockroach strain locate the left (Figure 2). This means that the wild cockroach strains were more sensitive to lambda-cyhalothrin, compared to the susceptible strain. This fact is also confirmed by observing a significant difference between treated lambda-cyhalothrin against the German cockroach strains compared to the other insecticides. It may be due to negative cross-resistance effect which occurs when a novel allele confers resistance to one toxic chemical or environmental factor and hyper-susceptibility to another.<sup>(43)</sup> That is to say, a trade-off occurs in which the benefit of resistance to one factor arising at the cost of greater susceptibility to a second factor.<sup>(44-46)</sup> Sequential deployment of negative cross-resistance insecticides is useful for insect control in few situations.<sup>(43)</sup>

Resistance level of the German cockroaches to the carbamates (propoxur), organophosphates (chlorpyrifos), pyrethroids (beta-cyfluthrin, deltamethrin), and phenyl pyrazole (fipronil) has increased to higher levels even in field populations. However, moderate level of resistance is found in German cockroaches against neonicotinoids (imidacloprid) and the oxadiazines like indoxacarb.<sup>(47,48)</sup> Currently, the German cockroaches are being emerged resistant to a wide range of insecticides including organophosphates, carbamates, pyrethroids, phenyl pyrazoles, spinosad, neonicotinoids and oxadiazines.<sup>(7,23-25,28-30,48)</sup>

The study supports that it had better not use the malathion and propoxur insecticides for control of the German cockroaches in these circumstances to prevent insecticide resistance or developing management. An appropriate choice is to use baits, particularly gel baits.<sup>(8,26,27,48)</sup> The other appropriate choice is to use lambda-cyhalothrin due to negative cross-resistance effect which occurs here.

## Conclusion

Like results of the study, it is well documented that the German cockroach is resistant to malathion and propoxur, while due to development of negative cross-resistance to lambda-cyhalothrin which occurs

here, the German cockroach is susceptible to lambda-cyhalothrin. The study supports that it had better not use the malathion and propoxur insecticides for control of the German cockroaches in these circumstances for preventing, developing and managing of the insecticide resistance. An appropriate choice is to use baits particularly gel baits. The other appropriate choice is to use lambda-cyhalothrin due to negative cross-resistance effect. This study provides a baseline of resistance data towards insecticides in German cockroach populations in Iran. The  $RR_{50}$  of the malathion and propoxur insecticides were about 5.0 folds. It is periodically a substantial need to evaluate sensitivity level of German cockroaches to consuming insecticides for predicting, preventing and managing of the insecticide resistance.

#### Acknowledgements

This study is the second author's M.Sc. thesis in Medical Entomology and Vector Control from the Department of Medical Entomology, School of Medicine, financially supported by Research Deputy of Hamadan University of Medical Sciences (Grant no: 9412187345).

#### Conflict of interest

We declare, hereby, that we have no conflict of interest.

#### References

1. Noureldin EM, Farrag HA. Population density of cockroach species and magnitude of their infestation in Jeddah Province, Saudi Arabia. *Flora and Fauna (jhansi)* 2008;2:259-67.
2. Paksa A, Ladonni H, Nasirian H. Comparison of PBO and DEF effects on creating bendiocarb and carbaryl insecticide resistance in German cockroach. *Sci J Kurdistan Univ Med Sci* 2012;17:91-101.
3. Nazari M, Motlagh BA, Nasirian H. Toxicity of cypermethrin and chlorpyrifos against German cockroach [*Blattella germanica* (Blattaria: Blattellidae)] strains from Hamadan, Iran. *Pak J Biol Sci* 2016;19: 259-64.
4. Tang QY, Jiang H, Li Y, Bourguignon T, Evans TA. Population structure of the German cockroach, *Blattella germanica*, shows two expansions across China. *Biological Invasions* 2016;18:2391-402.
5. Davari B, Hassanvand AE, Nasirian H, Ghiasian SA, Salehzadeh A, Nazari M. Comparison of cockroach fungal contamination in the clinical and non-clinical environments from Iran. *J Entomol Acarol Res* 2017;49: 109-15.
6. Nasirian H. Infestation of cockroaches (Insecta: Blattaria) in the human dwelling environments: A systematic review and meta-analysis. *Acta Trop* 2017; 167:86-98.
7. Sanei Dehkordi A, Salim Abadi Y, Nasirian H, Hazratian T, Gorouhi MA, Yousefi S, et al. Synergists action of piperonyl butoxide and S,S,S-tributyl phosphorotrithioate on toxicity of carbamate insecticides against *Blattella germanica*. *Asian Pac J Trop Med* 2017;10:981-6.
8. Davari B, Kashani S, Nasirian H, Nazari M, Salehzadeh A. Efficacy of Maxforce and Avion gel baits containing fipronil, clothianidin, and indoxacarb against the German cockroach (*Blattella germanica*): Gel bait efficacy against German cockroach. *Entomol Res* 2017;48:459-65.
9. Nasirian H, Salehzadeh A. Control of cockroaches (Blattaria) in sewers: a practical approach systematic review. *J Med Entomol* 2019;56:181-91.
10. Loucif L, Gacemi-Kirane D, Cherak Z, Chamlal N, Grainat N, Rolain JM. First report of German cockroaches (*Blattella germanica*) as reservoirs of CTX-M-15 extended-spectrum-beta-lactamase- and OXA-48 carbapenemase-producing Enterobacteriaceae in Batna University Hospital, Algeria. *Antimicrob Agents Chemother* 2016;60:6377-80.
11. Solomon F, Belayneh F, Kibru G, Ali S. Vector potential of *Blattella germanica* (L.) (Dictyoptera: Blattellidae) for medically important bacteria at food handling establishments in Jimma town, Southwest Ethiopia. *BioMed Res Int* 2016;2016:1-6.
12. Arif S, Taj M, Kamran K, Iqbal A, Mohammad G, Hussain K, et al. Household cockroaches of Quetta city as reservoir for infectious pathogenic bacteria. *J Entomol Zool Stud* 2017;5:649-53.
13. Martínez-Girón R, Martínez-Torre C, van Woerden HC. The prevalence of protozoa in the gut of German cockroaches (*Blattella germanica*) with special reference to *Lophomonas blattarum*. *Parasitol Res* 2017;116:3205-10.
14. Atiokeng Tatang RJ, Tsila HG, Wabo Pone J. Medically important parasites carried by cockroaches in Melong Subdivision, Littoral, Cameroon. *J Parasitol Res* 2017; 2017:7967325.
15. Nasirian H. Recent cockroach bacterial contamination trend in the human dwelling environments: A systematic review and meta-analysis. *Bangladesh J Med Sci* 2019;18:540-5.

16. Nasirian H. Contamination of cockroaches (Insecta: Blattaria) by medically important bacteria: a systematic review and meta-analysis. *J Med Entomol* 2019;56: 1534-54.
17. Nasirian H. Contamination of cockroaches (Insecta: Blattaria) to medically fungi: a systematic review and meta-analysis. *J Mycol Med* 2017;27:427-48.
18. Nasirian H. New aspects about *Supella longipalpa* (Blattaria: Blattellidae). *Asian Pac J Trop Biomed* 2016; 6:1065-75.
19. Ghavami M, Mohamadi J, Ladonni H. German cockroach resistance to permethrin in hospital, didactic therapeutic Shahid Beheshti Hospital in Zanjan. *J Zanjan Univ Med Sci* 2005;13:8-16.
20. Haghi F, Karami M, Enayati H, Safari R. Survey of susceptibility of cockroaches to pyrethroids insecticides in Sari hospital in 2004. *J Mazandaran Univ Med Sci* 2007;17:125-8.
21. Limoe M, Davari B, Moosa-Kazemi SH. Toxicity of Pyrethroid and Organophosphorous Insecticides against Two Field Collected Strains of the German Cockroach *Blattella germanica* (Blattaria: Blattellidae). *J Arthropod Borne Dis* 2012;6:112-8.
22. Ko AE, Bieman DN, Schal C, Silverman J. Insecticide resistance and diminished secondary kill performance of bait formulations against German cockroaches (Dictyoptera: Blattellidae). *Pest Manag Sci* 2016;72: 1778-84.
23. Jang CW, Ju Y, Chang KS. Insecticide Susceptibility of Field collected *Blattella germanica* (Blattaria: Blattellidae) in Busan, Republic of Korea during 2014. *Entomol Res* 2017;18: 2391-402.
24. Nasirian H, Ladoni H, Davari B, Shayeghi M, Yaghoobi Ershadi MR, Vatandoost H. Effect of fipronil on permethrin sensitive and permethrin resistant strains of *Blattella germanica*. *Sci J Kurdistan Univ Med Sci* 2006;11:33-41.
25. Nasirian H, Ladonni H, Vatandoost H. Duration of fipronil topical application toxicity in *Blattella germanica* field population strains. *Pakistan J Biol Sci* 2006;9:800-4.
26. Nasirian H. Duration of fipronil and imidacloprid gel baits toxicity against *Blattella germanica* strains of Iran. *Iran J Arthropod Borne Dis* 2007;1:40-7.
27. Nasirian H. Rapid elimination of German Cockroach, *Blattella germanica*, by fipronil and imidacloprid gel baits. *J Arthropod Borne Dis* 2008;2:37-43.
28. Nasirian H, Ladonni H, Shayeghi M, Ahmadi MS. Iranian non-responding contact method German cockroach permethrin resistance strains resulting from field pressure pyrethroid spraying. *Pak J Biol Sci* 2009;12:643-7.
29. Limoe M, Shayeghi M, Heidari J, Nasirian H, Ladonni H. Susceptibility level of hospital-collected strains of German cockroach *Blattella germanica* (L.) to carbamate and organophosphorous insecticides using surface contact method in Tehran (2007-2008). *J Kermanshah Univ Med Sci* 2010;13:337-43.
30. Ladonni H, Paksa A, Nasirian H, Doroudgar A, Abaie M. Detection of Carbamate and organo phosphorus susceptibility levels in German cockroach in vivo. *Toloo e Behdasht* 2013;40:95-105.
31. Abd-Elghafar S, Appel A, Mack T. Toxicity of several insecticide formulations against adult German cockroaches (Dictyoptera: Blattellidae). *J Econ Entomol* 1990;83:2290-4.
32. Diaz Pantoja C, Perez MG, Calvo E, Rodriguez MM, Bisset JA. Insecticide resistance studies on *Blattella germanica* (Dictyoptera: Blattellidae) from Cuba. *Ann NY Acad Sci* 2000;916:628-34.
33. Diaz C, Enriquez D, Bisset JA. Status of resistance to insecticides in field strains of the *Blattella germanica* species (Dictyoptera: Blattellidae) from Pinar del Rio municipality. *Rev Cubana Med Trop* 2003;55:196-202.
34. Hemingway J, Small GJ, Monro AG. Possible mechanisms of organophosphorus and carbamate insecticide resistance in German cockroaches (Dictyoptera: Blattellidae) from different geographical areas. *J Econ Entomol* 1993;86:1623-30.
35. Cochran D. Looking into cockroach resistance extensive research on insecticide resistance in German cockroach yields surprising results. *Pest Management* 1995;3:31-5.
36. Gliniewicz A, Krzemińska A, Sawicka B. Susceptibility of cockroaches *Blattella germanica* L. collected from hospitals to selected pyrethroid and carbamate insecticides. *Rocz Panstw Zakl Hig* 1996;47:333-41.
37. Lee C, Yap H, Chong N, Lee R. Insecticide resistance and synergism in field collected German cockroaches (Dictyoptera: Blattellidae) in Peninsular Malaysia. *Bull Entomol Res* 1996;86:675-82.
38. Diaz C, Enriquez D, Bisset J. Status of resistance to insecticides in field strains of the *Blattella germanica* species (Dictyoptera: Blattellidae) from Pinar del Rio municipality. *Rev Cubana Med Trop* 2003;55:196-202.
39. Pai HH, Wu SC, Hsu EL. Insecticide resistance in German cockroaches (*Blattella germanica*) from hospitals and households in Taiwan. *Int J Environ Health Res* 2005;15:33-40.
40. Wu X, Appel AG. Insecticide Resistance of Several



- Field-Collected German Cockroach (Dictyoptera: Blattellidae) Strains. *J Econ Entomol* 2017;110:1203-9.
41. Diaz PC, Alvarez Gavilan Y, de Armas Rodriguez Y, Bisset Lazcano JA. Determination of insecticide-resistance and resistance mechanisms of *Blattella germanica* (Dictyoptera: Blattellidae). *Rev Cubana Med Trop* 2007;59:159-65.
  42. Fardisi M, Gondhalekar AD, Scharf ME. Development of diagnostic insecticide concentrations and assessment of insecticide susceptibility in German cockroach (Dictyoptera: Blattellidae) field strains collected from public housing. *J Econ Entomol* 2017; 110:1210-7.
  43. Pittendrigh B, Gaffney P. Pesticide resistance: can we make it a renewable resource? *J Theor Biol* 2001;211: 365-75.
  44. Scharf ME, Kaakeh W, Bennett GW. Changes in an insecticide-resistant field population of German cockroach (Dictyoptera: Blattellidae) after exposure to an insecticide mixture. *J Econ Entomol* 1997;90: 38-48.
  45. Pittendrigh BR, Gaffney P, Murdock LL. Deterministic modeling of negative cross-resistance strategies for use in transgenic host-plant resistance. *J Theor Biol* 2000;204:135-50.
  46. Kolaczinski JH, Curtis CF. Investigation of negative cross-resistance as a resistance-management tool for insecticide-treated nets. *J Med Entomol* 2004;41: 930-4.
  47. Chai RY, Lee CY. Insecticide resistance profiles and synergism in field populations of the German cockroach (Dictyoptera: Blattellidae) from Singapore. *J Econ Entomol* 2010;103:460-71.
  48. Naqqash MN, Gökçe A, Bakhsh A, Salim M. Insecticide resistance and its molecular basis in urban insect pests. *Parasitol Res* 2016;115:1363-73.